

DOE/ET/53088-6

IFSR #6

STATISTICAL CHARACTERIZATION
OF
PERIODIC, MEASURE-PRESERVING MAPPINGS

John R. Cary, James D. Meiss
and
A. Bhattacharjee

Institute for Fusion Studies
The University of Texas at Austin
Austin, Texas 78712

January, 1981

STATISTICAL CHARACTERIZATION
OF
PERIODIC, MEASURE-PRESERVING MAPPINGS

John R. Cary, James D. Meiss
and
A. Bhattacharjee

Institute for Fusion Studies
University of Texas
Austin, Texas 78712
January, 1981

ABSTRACT

A method of statistically characterizing an area-preserving, doubly periodic mapping is presented. This method allows one to calculate the characteristic functions, the diffusion coefficient, the correlation functions, and the joint probabilities of the mapping.

The standard mapping (Ref. 1, Sec. 5) has been studied extensively as a model of stochastic behavior. Early work included calculations of the perturbed invariants for small values of the nonlinearity parameter and the study of the transition to stochasticity². More recently, there has been considerable interest in the study of the statistical description of the stochastic regime with the calculation of the diffusion coefficient^{3,4}. Here we present a method for statistically characterizing the standard mapping,

$$\begin{aligned} p_{n+1} &= p_n + \varepsilon \sin(x_n) \\ x_{n+1} &= x_n + p_{n+1}, \end{aligned} \quad (1)$$

and other doubly periodic mappings (Ref. 1, Sec. 5.1). The essence of our method is the calculation of the characteristic functions of the mapping. These characteristic functions allow us to determine various statistical quantities, such as the diffusion coefficient, the correlation functions, and the joint probability distributions.

The characteristic functions are defined with integer argument as follows:

$$\begin{aligned} \chi_0(m_0) &\equiv \langle \exp(im_0 x_n) \rangle \\ \chi_1(m_0, m_1) &\equiv \langle \exp(im_0 x_n + im_1 x_{n+1}) \rangle \\ \chi_2(m_0, m_1, m_2) &\equiv \langle \exp(im_0 x_n + im_1 x_{n+1} + im_2 x_{n+2}) \rangle \end{aligned} \quad (2)$$

The angular brackets in these definitions are left unspecified for the moment, except that they are time-translation invariant. An example of a time-translation invariant average is the average of the initial conditions over an invariant region R of phase space⁵, i.e. a region which is mapped onto itself:

$$\begin{aligned} \langle \exp(im_0 x_n + im_1 x_{n+1} + im_2 x_{n+2}) \rangle &\equiv \left(\int_R dp_0 dx_0 \right)^{-1} \\ &\times \int_R dp_0 dx_0 \exp[im_0 x_n(x_0, p_0) + im_1 x_{n+1}(x_0, p_0) + im_2 x_{n+2}(x_0, p_0)] \end{aligned} \quad (3)$$

The crucial point of this analysis is that the mapping equations (1) in the form

$$x_{n+k} = 2x_{n+k-1} - x_{n+k-2} + \varepsilon \sin(x_{n+k-1}) \quad (4)$$

allow one to express x_{n+1} in terms of x_n for $n \geq 1$. By using the Bessel function identity $\exp(i\varepsilon \sin(x)) = J_\ell(\varepsilon) \exp(i\ell x)$, we find

$$\begin{aligned} \chi_k(m_0, m_1, \dots, m_k) &\equiv \langle \exp(im_0 x_n + im_1 x_{n+1} + im_2 x_{n+2} + \dots + im_k x_{n+k}) \rangle \\ &\equiv \langle \exp[im_0 x_n + \dots + im_{k-3} x_{n+k-3} + i(m_{k-2} - m_k) x_{n+k-2} \\ &\quad + i(m_{k-1} + 2m_k) x_{n+k-1} + im_k \varepsilon \sin(x_{n+k-1})] \rangle \\ &= \sum_{\ell} J_{\ell}(m_k \varepsilon) \chi_{k-1}(m_0, \dots, m_{k-3}, m_{k-2} - m_k, m_{k-1} + 2m_k + \ell). \end{aligned} \quad (5)$$

Moreover, χ_1 itself may be calculated by Eq. (3):

$$\chi_1(m_0, m_1) = \left(\int_R dp_0 dx_0 \right)^{-1} \int_R dp_0 dx_0 \exp[i(m_0 + m_1)x_0 - im_0 p_0]. \quad (6)$$

The interpretation of this method so that we must first select a class of orbits and, thereby, an invariant region R. Performing the integrals in Eq. (6) we obtain the characteristic functions averaged over that class of orbits. All subsequent quantities derived from the characteristics functions are therefore averages over that class of orbits.

Let us take the region R in Eq. (6) to be all of phase space. Eqs. (5) and (6) immediately yield

$$\chi_1(m_0, m_1) = \delta_{m_0, 0} \delta_{m_1, 0}, \quad (7a)$$

$$\chi_2(m_0, m_1, m_2) = \delta_{m_0, m_2} J_{-m_1 - 2m_0}^{(m_2 \epsilon)} \quad (7b)$$

$$\text{and } \chi_3(m_0, m_1, m_2, m_3) = J_{m_3 - m_2 - 2m_0}^{(m_0 \epsilon)} J_{m_0 - m_2 - 2m_3}^{(m_3 \epsilon)} \quad (7c)$$

Knowledge of the characteristic functions allows us to calculate the averaged diffusion constant,

$$D \equiv \left\langle \lim_{T \rightarrow \infty} \frac{1}{2T} (p_f - p_i)^2 \right\rangle = \left\langle \lim_{T \rightarrow \infty} \frac{1}{2T} \sum_{i=1}^T \sum_{j=1}^T \Delta p_i \Delta p_j \right\rangle. \quad (8)$$

To reduce the double sum in the last equation, we rewrite it in the form,

$$D = \lim_{T \rightarrow \infty} \frac{1}{2T} \sum_{i=1}^T \sum_{j=1-i}^{T-i} \langle \Delta p_i \Delta p_{i+j} \rangle. \quad (9)$$

Next we assume that the terms in the sum fall off rapidly with j , so that as $T \rightarrow \infty$, we can sum over all j .

$$D = \lim_{T \rightarrow \infty} \frac{1}{2T} \sum_{i=1}^T \sum_{j=-\infty}^{\infty} \langle \Delta p_i \Delta p_{i+j} \rangle. \quad (10)$$

Then we use the time translation invariance of the average to sum over i :

$$D = \frac{1}{2} \sum_{j=-\infty}^{\infty} \langle \Delta p_i \Delta p_{i+j} \rangle. \quad (11)$$

Finally, we use the Fourier expansion, $\Delta p_j = \epsilon \sin(x_j) = (-i/2) [\exp(ix_j) - \exp(-ix_j)]$, to obtain the following formula for the diffusion coefficient,

$$D = \frac{1}{4} \epsilon^2 \operatorname{Re} \left\{ \chi_0(0) - \chi_0(2) + 2 \sum_{j=1}^{\infty} [\chi_j(1,0,\dots,0,-1) - \chi_j(1,0,\dots,0,1)] \right\}. \quad (12)$$

Inserting the special solution of Eqs. (7), we find

$$\begin{aligned}
D = \frac{1}{4} \varepsilon^2 \left\{ & 1 - 2J_2(\varepsilon) + 2J_3^2(\varepsilon) - 2J_1^2(\varepsilon) + 2 \sum_{n=-\infty}^{\infty} J_n(\varepsilon) J_{-n}(\varepsilon) J_{-2(n+2)} [(n+2)\varepsilon] \right. \\
& - 2 \sum_{n=-\infty}^{\infty} J_n^2(\varepsilon) J_{-2(n+1)} [(n+2)\varepsilon] \\
& \left. + 2 \sum_{j=5}^{\infty} [\chi_j(1,0,0,\dots,0,\pm 1) - \chi_j(1,0,0,\dots,0,1)] \right\} \quad (13)
\end{aligned}$$

We note that the first four terms of this series are exactly those found by Rechester and White³. (We also note that Karney et al.⁶ found those same terms by computing the force correlation functions.)

A knowledge of the characteristic functions allows one also to calculate the correlation functions,

$$C_m \equiv \langle \text{fr}(x_n) \text{fr}(x_{n+m}) \rangle, \quad (14)$$

where $\text{fr}(x)$ is the number between zero and 2π which equals x modulo 2π . The function $\text{fr}(x)$ can be written as a Fourier series,

$$\text{fr}(x) = \sum_{k=-\infty}^{\infty} r_k e^{ikx}, \quad (15)$$

where $r_0 = \pi$ and $r_k = i/k$ for $k \neq 0$. By straightforward algebraic manipulation we find

$$C_m = \pi^2 + \operatorname{Re} \left[4\pi \sum_{k=1}^{\infty} \frac{i\chi_0(k)}{k} + 2 \sum_{k=1}^{\infty} \sum_{\ell=1}^{\infty} \frac{\chi_m(k, 0, 0, \dots, 0, -\ell) - \chi_m(k, 0, 0, \dots, 0, 2)}{\ell k} \right] \quad (16)$$

For the solution of Eq. (7) we obtain

$$C_0 = \frac{4}{3} \pi^2 \quad (17a)$$

$$C_1 = \pi^2 \quad (17b)$$

$$C_2 = \pi^2 - 2 \sum_{k=1}^{\infty} \frac{J_{2k}(k\varepsilon)}{k^2} \quad (17c)$$

$$C_3 = \pi^2 - 2 \sum_{k=1}^{\infty} \sum_{\ell=1}^{\infty} \frac{J_{\ell-2k}(k\varepsilon) J_{k-2\ell}(\ell\varepsilon) - (-1)^{\ell+k} J_{\ell+2k}(k\varepsilon) J_{k+2\ell}(\ell\varepsilon)}{\ell k} \quad (17d)$$

It should be noted that we have made an error in our application of this method to the calculation of the diffusion coefficient by neglecting the accelerator modes (Ref. 11, Sec. 5.5; Ref. 6). Accelerator modes exist throughout a region of finite measure in phase space. They have the property that p_n continuously increases or decreases roughly as n . Such modes contribute an infinite diffusion when folded into the average (8) and will cause the series (13) not to converge.

To correct for the presence of accelerator modes, we should

choose the region R of Eqs. (3) and (6) to be the stochastic region of phase space. Our value for $\chi_1(0,0)$ remains unchanged by this choice, but now $\chi_1(m,n)$ for m or n not equal to zero no longer vanishes. Instead it is of the order of $1/\epsilon^2$ since the relative measure of the accelerator regions is of that order (Ref. 1, Sec. 5.5). This will cause corrections to Eq. (13) of order equal to that of the last sum.

To illustrate the generality of this method, we consider its application to the mapping,

$$u_{j+1} = u_j + \mu - \nu \cos v_j$$

$$v_{j+1} = v_j + \mu + \nu \sin u_{j+1} ,$$

which has been discussed by Karney⁷. If we define the characteristic functions to be

$$\chi_{M,N}(m_0, \dots, m_M, n_0, \dots, n_N) \equiv \langle \exp(im_0 u_j + im_1 u_{j+1} + \dots + im_M u_{j+M} + in_0 v_j + in_1 v_{j+1} + \dots + in_N v_{j+N}) \rangle ,$$

we immediately find the recursion relations,

$$\chi_{M,M} = \sum_{\ell} J_{\ell}(n_M \nu) e^{in_M \mu} \chi_{M,M-1}(m_0, m_1, \dots, m_{M-1}, m_M + \ell, n_0, n_1, \dots, n_{M-2}, n_{M-1} + n_M)$$

and

$$\chi_{M,M-1} = \sum_{\ell} J_{\ell}(m_M, \nu) e^{i(m_M \ell - \ell \pi/2)} \chi_{M-1,M-1}(m_0, \dots, m_{M-1} + m_M, n_0, \dots, m_{M-1} + \ell).$$

Finally, we would like to note that this technique allows one to calculate the fractionally reduced joint probability distribution $P(\bar{x}_0, \bar{p}_0, \bar{x}_k, \bar{p}_k)$ which has been discussed by Grebogi et al.⁸ For example

$$\begin{aligned} P(\bar{x}_0, \bar{p}_0, \bar{x}_k, \bar{p}_k) &= \langle \delta[\bar{x}_0 - \text{fr}(x_j)] \delta[\bar{p}_0 - \text{fr}(p_j)] \delta[\bar{x}_k - \text{fr}(x_{j+k})] \\ &\quad \delta[\bar{p}_k - \text{fr}(p_{j+k})] \rangle \\ &= \left(\frac{1}{2\pi}\right)^4 \sum_{m_0, m_k, n_0, n_k} \exp(-im_0 \bar{x}_0 - im_k \bar{x}_k - in_0 \bar{p}_0 - in_k \bar{p}_k) \\ &\quad \times \chi_{k,k}(m_0, 0, 0, \dots, m_k, n_0, 0, 0, \dots, n_k). \end{aligned}$$

Obviously this method can be used to find the characteristic functions of any doubly periodic mapping. Furthermore, in the stochastic regime, where the higher order characteristics are small, one can use this method to obtain an approximate expression for the diffusion coefficient, the correlation functions, and the joint probabilities.

References

1. Boris V. Chirikov, *Physics Reports* 52, 263 (1979).
2. J.M. Greene, *J. Math. Phys.* 20, 1183 (1979).
3. A.B. Rechester and R.B. White, *Phys. Rev. Lett.* 44, 1586 (1980).
4. A.B. Rechester, M.N. Rosenbluth, and R.B. White, *Plasma Physics Rep. PPPL-1678*, Princeton University (1980).
5. V.I. Arnold and A. Avez, *Ergodic Problems of Classical Mechanics*, (W.A. Benjamin, Inc., New York, 1968) p.15-19.
6. C.F.F. Karney, A.B. Rechester, and R.B. White, *Plasma Physics Rep. PPL-1752*, Princeton University (1981).
7. Charles F.F. Karney, *Phys. Fluids* 22, 2188 (1979).
8. Celso Grebogi, Allan N. Kaufman, and Henry D.I. Abarbanel, *Bull. Am. Phys. Soc.* 25, 988 (1980).